



Anti-T. pallidum Polyclonal antibody (DPAB0151)

This product is for research use only and is not intended for diagnostic use.

PRODUCT INFORMATION

Specificity	Recognizes all antigens. Non-reactive in RPR assay. This antibody has not been absorbed and may react with related microorganisms.
Target	T. pallidum
Immunogen	Highly purified T. pallidum
Source/Host	Rabbit
Species Reactivity	T. pallidum
Purification	Purified IgG fraction covalently coupled with high purity Isomer I of fluorescein isothiocyanate. Care is taken to ensure complete removal of any free fluorescein from the final product
Conjugate	FITC
Applications	Suitable for use in direct IFA and immunohistochemistry (paraffin). Applications include direct FA staining of target antigens in a permissive tissue culture system. Acetone fixation of the antigen source is recommended prior to staining. Each laboratory should determine an optimum working titer for use in its particular application. Other applications have not been tested but use in such assays should not necessarily be excluded.
Applications Concentration	FA staining of target antigens in a permissive tissue culture system. Acetone fixation of the antigen source is recommended prior to staining. Each laboratory should determine an optimum working titer for use in its particular application. Other applications have not been
	FA staining of target antigens in a permissive tissue culture system. Acetone fixation of the antigen source is recommended prior to staining. Each laboratory should determine an optimum working titer for use in its particular application. Other applications have not been tested but use in such assays should not necessarily be excluded.
Concentration	FA staining of target antigens in a permissive tissue culture system. Acetone fixation of the antigen source is recommended prior to staining. Each laboratory should determine an optimum working titer for use in its particular application. Other applications have not been tested but use in such assays should not necessarily be excluded. 4-5mg/ml (OD280nm, E0.1% = 1.4)
Concentration	FA staining of target antigens in a permissive tissue culture system. Acetone fixation of the antigen source is recommended prior to staining. Each laboratory should determine an optimum working titer for use in its particular application. Other applications have not been tested but use in such assays should not necessarily be excluded. 4-5mg/ml (OD280nm, E0.1% = 1.4) 1 ml

45-1 Ramsey Road, Shirley, NY 11967, USA

Email: info@creative-diagnostics.com

Tel: 1-631-624-4882 Fax: 1-631-938-8221

BACKGROUND

Introduction	Treponema pallidum is a species of spirochaete bacterium with subspecies that cause treponemal diseases such as syphilis, bejel, pinta and yaws. The treponemes have a cytoplasmic and outer membrane. They are not seen on a Gram stained smear because the organism is too thin to be observed under a light microscope.
Keywords	Bacteria; Spirochaetes; Spirochaetales; Spirochaetaceae; Treponema; T. pallidum; Treponema pallidum